

# Discrete Markov chains and mixing times

## Homework 1

Due 3 月 4 日 (周一) at the start of class

### 1 Textbook problems

- Chapter 1: 1.5, 1.6, 1.10, 1.11, 1.12
- Chapter 2: 2.3, 2.5 (give two different proofs), 2.6, 2.7

Other exercises in chapters 1 and 2 may also be interesting but only the above problems are required. Others exercises are suggested for interested students for additional review and learning.

### 2 Additional problems

As in class, let  $\mathbb{M}_n$  be the collection of  $n \times n$  stochastic matrices. Let  $\mathbb{M}_n^I$  be those which are irreducible. Assume  $|\mathcal{X}| < \infty$ .

1. Classify the elements of  $\mathbb{M}_n$  for which the uniform distribution  $\pi = (1/n, \dots, 1/n)$  is stationary. What kind of graphs satisfy this condition?
2. At the start of the proof of Proposition 1.14, we used the equivalence of  $\mathbb{P}_z(\tau_z^+ < \infty) = 1$  and  $\mathbb{E}_z(\tau_z^+) < \infty$  (which we proved in §1.7). Where else did the proof use the assumption  $\mathbb{P}_z(\tau_z^+ < \infty) = 1$ ?
3. We stated in class that  $x \in \mathcal{X}$  is essential iff  $\mathbb{P}_x(\tau_x^+ < \infty) = 1$ . Prove this.

### 3 Optional problems

Interested students may consider the following problem. It is not required. Let  $\mathcal{P}(\mathcal{X})$  be the collection of probability measures on  $\mathcal{X}$ .

1. (a) Consider the map  $S : \mathbb{M}_n^I \rightarrow \mathcal{P}(\mathcal{X})$  given by  $S(P) = \pi$ , where  $\pi$  is the (unique) stationary distribution of  $P$ . Is  $S$  injective? Is  $S$  surjective?
- (b) Consider the reversal map  $\hat{\cdot} : \mathbb{M}_n^I \rightarrow \mathbb{M}_n^I$  given by  $\hat{\cdot}(P) = \hat{P}$ . It is easy to see that  $\hat{\cdot}$  is a bijection. Does it possess other nice operator-theoretic properties? (For example, is it an isometry, in some sense? Does it preserve the spectrum?)